

Spectral Properties of Heavy Ions Associated with Interplanetary Shocks at 1 AU

**SHINE 2004
Big Sky, Montana**

M. I. Desai

University of Maryland, College Park, MD 20742, USA

Co-Authors:

G. M. Mason: University of Maryland

C.M.S.Cohen & M. E. Wiedenbeck: SRL, Caltech, CA

J. E. Mazur: Aerospace Corp

J. R. Dwyer: Florida Institute of Technology



R.E. Gold & S. M. Krimigis: JHU/APL

C.W.Smith: University of New Hampshire

Q. Hu: IGPP, UC Riverside, CA

R. M. Skoug: Los Alamos National Laboratory


CME-driven Interplanetary Shocks



Intensity-time profiles for an IP shock


- Surveyed 72 shocks between Oct. 1997-Oct. 2002
- ^3He ions accelerated in 45 IP shocks
- Pre-shock intervals provide a "proxy" for ambient particles in the IP medium

(Desai et al. 2001, *ApJ* 553, L89 & 2003, *ApJ*, 588, 1149).




Ambient and SEP Abundances

*Ambient material
comprises ~30%
from impulsive
flares, and ~70%
from large
gradual SEPs*



(Desai et al. 2003 ApJ, 588, 1149).

IP Shock compared with solar wind abundances




- *Shock abundances are poorly correlated with solar wind abundances*
- *No clear dependence on M/Q*
- *Difficult to understand in terms of rigidity-dependent acceleration of solar wind ions*

(Desai et al. 2003 ApJ, 588, 1149).

IP Shock compared with ambient suprathermal abundances


- *Shock abundances are well correlated with ambient suprathermal abundances*
- *Exhibit a M/Q -dependent depletion*
- *Consistent with rigidity-dependent shock acceleration of ambient suprathermals*



(Desai et al. 2003 ApJ, 588, 1149).

Fe/O at IP shocks is depleted relative to ambient values

Larger decrease at higher energy




(Desai et al. To appear in ApJ. Aug. 20, 2004 issue)

Energy Spectra during an IP shock:

All Spectra fitted by $j = j_0 E^\alpha \exp(-E/E_0)$



however, α and E_0 are coupled;
Use only 0.1-0.5 MeV n⁻¹ to obtain the power-law indices



Parameter	Carbon	Oxygen	Iron
No. of Points	8	10	8
j_0	4.64 ± 0.87	9.32 ± 1.21	3.17 ± 1.49
α	1.36 ± 0.11	1.33 ± 0.09	1.08 ± 0.27
E_0	0.69 ± 0.09	0.72 ± 0.06	0.35 ± 0.06
α^2/α	0.87	0.91	0.88

Spectral indices of C, O, & Fe



Differences in Fe and O indices are at odds with injection of a mono-energetic seed population



(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

O e-folding energy vs. shock parameters

E-folding energy is independent of local shock parameters




(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)


2-hr. av. O spectral index vs. $(M+2)/(2M-2)$

- Spectral index is poorly correlated with compression ratio, M
- Results are at odds with predictions of simple steady-state models

(Desai et al. To appear in *ApJ*. Aug. 20, 2004 issue)





IP shock event measured by ULEIS & SIS




(Desai et al. To appear in *ApJ*. Aug. 20, 2004 issue)

Energy spectra measured by ULEIS & SIS




(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

3 Classes of Fe/O energy-dependence



(Desai et al. To appear in *ApJ*. Aug. 20, 2004 issue)


Energy-dependence of Fe/O



$$\square_{\text{Fe}} = \frac{\text{Fe/O (0.62 MeV/n.)}}{\text{Fe/O (0.22 MeV/n.)}}$$


(Desai et al. To appear in *ApJ*. Aug. 20, 2004 issue)

Γ_{Fe} vs. $^3\text{He}/^4\text{He}$ ratio; $^3\text{He}/^4\text{He}$ ratio vs. θ_{Bn}
Extreme Events Only



(Adapted from Desai et al. 2004, To appear in *ApJ*. Aug. 20, 2004)

Γ_{Fe} vs. $^3\text{He}/^4\text{He}$ ratio; $^3\text{He}/^4\text{He}$ ratio vs. θ_{Bn}
All events




(Desai et al. To appear in *ApJ*. Aug. 20, 2004 issue)

$^3\text{He}/^4\text{He}$ ratio vs. Injection Threshold Speed, $V_{inj} = V_s \cdot \sec(\theta_{Bn})$



Most IP shocks including the 3 with rising Fe/O ratios have $V_{inj} < 2 \cdot V_{sw}$

$^3\text{He}/^4\text{He}$ ratio and \square_{Fe} are poorly correlated with V_{inj}

Conclusion: Injection threshold speeds do not appear to play a significant role in the energy-dependent behavior of Fe/O




Γ_{Fe} vs. O spectral index & O fluence



(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

Fe/O at IP shocks vs. ambient


*Fe/O ratios at IP shocks
and ambient are well
correlated*



(Desai et al. 2003 ApJ, 588, 1149).



Shock C/O & Fe/O normalized to ambient values

- *Fe/O ratios are depleted by ~30% relative to ambient values*
- *Energy-dependence of Fe/O is diminished when compared with ambient Fe/O - not expected from mono-energetic injection*



(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

O Spectral index & Fe/O dependence at IP shocks vs. Ambient




(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

Summary

- *Spectral parameters and energy-dependence of Fe/O are independent of local shock parameters*
- *5 out of 72 events (~7%) have rising Fe/O with energy; Fe/O in other events are constant or decrease with energy*
- *Fe/O at IP shocks are typically ~30% lower than in the ambient population*
- *The O spectra and energy-dependence of Fe/O are similar at IP shocks and in the ambient population*

Sketch of Re-acceleration of Seed spectra at IP shocks



(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

Conclusion

IP shocks accelerate seed spectra composed of suprathermal ions from gradual and impulsive SEP events by a systematic rigidity-dependent mechanism where ions with higher M/Q are accelerated less efficiently than those with lower M/Q


Relevant Issues

- *How common are IP shock events with rising Fe/O ratios?*
 - *~7% of events have rising Fe/O with energy*
- *What are the key differences between IP shocks with rising and decreasing Fe/O ratios?*
 - *No appreciable differences in shock properties*
- *Does any particular local shock parameter play a role in determining the energy-dependent behavior of Fe/O?*
 - *Cannot rule this out completely, but no evidence that local shock properties are important*
- *The primary cause of rising Fe/O in IP shocks*
 - *Re-acceleration of energetic ion seed spectra that themselves have rising Fe/O with energy*

THE END


He mass histograms during quiet periods 1997-1999

- *Energetic ^3He ions from impulsive flares are present in the interplanetary medium*
- *Remnant impulsive ^3He -rich material could be re-accelerated at CME-driven IP shocks*
- *IP shock abundances should exhibit solar cycle dependence*




(Mason et al. 1999 ApJ, 525, L133).

Wind/STEP ~ 100 keV n^{-1} Fe Intensity




Solar Wind Abundance Ratios

von Steig



Von Steiger et al., (2000, JGR , vol. 105, 27217)


Shock Abundances and Ambient Abundances



(Desai et al. 2003 ApJ, 588, 1149).


*Intensity-time profiles
for an IP shock*

*Surveyed 72 IP shocks
between Oct. 1997-
Oct. 2002*



(Desai et al. 2003 ApJ, 588, 1149).


Shock abundances versus solar wind



(Desai et al. 2003 ApJ, 588, 1149).


Solar cycle dependence of shock abundances

- *Fe/O ratio in interplanetary medium and in IP shocks show clear solar cycle dependence*
- *Solar wind abundances show no such solar cycle effects*
- *Dominant source of material is not solar wind but interplanetary suprathermals*



(Desai et al. 2003 ApJ, 588, 1149).


Frequency of ^3He -rich shocks, fraction of time ^3He ions are present in the IP medium



(from Desai et al. 2003, ApJ vol. 558, 1149; Wiedenbeck et al SW10).


Shock/Solar Wind Abundances versus M/Q

- *IP shock and solar wind abundances are poorly correlated*
- *Fe is enhanced, while C is depleted by ~40%*
- *Difficult to understand in terms of rigidity-dependent acceleration*



(Desai et al. 2003 ApJ, 588, 1149).

Peak Oxygen Intensity at shock vs. Oxygen Fluence during a 5-day interval upstream




Contributions to Long-Term O Fluences

Suprathermal tail comprises:

- *Solar wind suprathermals*
- *Interstellar and inner source pickup ions*
- *Residual suprathermals from solar and interplanetary activity*


(Mewaldt et al., 2001 AIP conference Proceedings 598, 165)



Wind/STEP ~ 100 keV n^{-1} C/O and Fe/O ratios


C/O and Fe/O ratios exhibit a clear solar cycle behavior

*CIR-like at solar minimum
SEP-like at solar maximum*





Shock /Ambient versus M/Q

- *C/O & N/O ratios similar in IP shocks and ambient*
- *Ne-Fe abundances show an M/Q-dependent depletion; Fe is depleted by 30-40%*
- *Systematic differences due to rigidity-dependent acceleration*



(Desai et al. 2003 ApJ, 588, 1149).







3 Classes of Fe/O energy-dependence



(Desai et al. *To appear in ApJ. Aug. 20, 2004 issue*)

Helium Mass histograms for 6 IP shocks


- ^3He ions accelerated in 45 IP shocks
- Enhancements between factors of ~ 4 -2000 over solar wind value
- $^3\text{He}/^4\text{He}$ ratio ~ 100 times greater the solar wind value in 7 events



(Desai et al. ApJ 2001 553, L89)

Rigidity-dependent acceleration at shocks

Iron is accelerated less efficiently than oxygen



Motivation

- *How common are IP shock events with rising Fe/O ratios?*
- *What are the key differences between IP shocks with rising and decreasing Fe/O ratios?*
- *Does any particular local shock parameter play a role in determining the energy-dependent behavior of Fe/O?*
- *What causes the energy-dependent behavior of Fe/O in IP shocks?*