

Heliospheric flux and ICMs

Mathew Owens



Center for space physics
Boston University



Overview

1. Background

- Heliospheric flux catastrophe
- $|B|$ with solar cycle – explained by ICMEs?

2. How much flux is injected by ICMEs?

- New flux-rope model of MCs

3. What “disconnection rate” is required to match observations?

- “Back of the envelope” calculation



Center for space physics
Boston University



Flux from CMEs

- CMEs involve the eruption of closed magnetic field systems
- At 1 AU BDEs suggest ICMEs are still (partially) closed structures, and thus add to the heliospheric flux budget
- ICMEs must eventually open to avoid the heliospheric “flux catastrophe” (*McComas et al., 1992*)




Center for space physics
Boston University





Heliospheric flux variation

- Solar cycle variation in $|B|$
- *Richardson et al. [2002]*
 - Only s/w in corotating streams carries solar cycle variation of $|B|$
 - S/W in **recognisable** ICMEs has different variation




How closed are ICMEs?


- Magnetic clouds at 1 AU are $\sim 50\%$ open
- Clouds at 5 AU \sim as closed as those at 1 AU
- Results imply
 - $\sim 50\%$ of fields open during CME liftoff
 - Fields must open over very long timescales




Rate of ICME opening

- Variation of closed nature of ICMEs with distance
 - No discernable trend
- Rate of opening below detection threshold
- Sampling same ICME at difference times
 - Closed fields increase?
(*Skoug et al.*, 2000)
- Sensitive to spacecraft trajectory through ICME?





- At CME liftoff
 - a. Partial disconnection (closed-closed) creates flux rope coil
 - b. Interchange reconnection (closed-open) opens coil
- In the following months to years
 - Interchange reconnection at Sun opens remaining field lines
- Meanwhile closed fields in legs of ICMEs far out in heliosphere may
 - no longer exhibit counterstreaming
 - be counted as open fields
 - cause solar cycle rise in $|B|$



What is the typical flux injected by a single CME?


- Typical dimming “foot-point” fluxes
 - E.g. May 12th 1997 (*Webb et al*, 2001) : 1×10^{21} Mx
- How much flux is seen at 1 AU?
 - Estimates made using force-free models
 - E.g. May 12th 1997 (*Webb et al*, 2001) : 7×10^{20} Mx
- Problems:
 - Axis orientation errors – MC thickness: $f(\cos[\theta_{YGSE}])$
 - Axial field strength (i.e. estimate of closest approach)
 - **Cross-sectional shape/extent**






Center for space physics
Boston University



New flux-rope model of MCs



Model validation




Center for space physics
Boston University



Flux estimates using new model

- Estimate for May 12th 1997:
 $1 \times 10^{21} \text{ Mx}$
 - However, can vary from force-free model by ~ 10
- *Lynch et al.* [2005] survey of 132 MCs using force free model:
 $\langle _ \rangle \sim 6 \times 10^{20} \text{ Mx}$
Spread over 3 orders of magnitude
- Fitting to **limited number** of MCs with **new model**:
 $\langle _ \rangle \sim 10^{21} \text{ Mx}$



Back of the envelope calculation

- $\varphi_H = \varphi_{CH} + \varphi_{CME}$
- Closed flux from a single CME:
 $\varphi = \varphi_0(1-D_0) - D(t)$
- Assume: $\frac{d\varphi}{dt} \propto \varphi$ i.e. $\varphi = \varphi_0(1-D_0) e^{-t}$
- Sum over all CMEs: $\Phi_{CME} = \sum_{n=1}^{\infty} \varphi_n = \sum_{n=1}^{\infty} \varphi_{0n}(1-D_{0n}) \exp[-\lambda_n t_n]$
- “Average” ICME: $\Phi_{CME} = \overline{\varphi_0} \left(1 - \overline{D_0}\right) \sum_{n=1}^{\infty} \exp[-\lambda_n / f]$

Back of the envelope calculation

- Solar min/max:

$$\Phi_H(MIN) = \pi(1AU)^2 |B_{MIN}| \cos(45^0) = \Phi_{CH}(MIN) + \overline{\varphi_0} (1 - \overline{D_0}) \sum_{n=1}^{\infty} \exp[-\lambda n / f_{MIN}]$$

$$\Phi_H(MAX) = \pi(1AU)^2 |B_{MAX}| \cos(45^0) = \Phi_{CH}(MAX) + \overline{\varphi_0} (1 - \overline{D_0}) \sum_{n=1}^{\infty} \exp[-\lambda n / f_{MAX}]$$

- Assume $\Phi_{CH}(MIN) = \Phi_{CH}(MAX)$ (i.e. all variation in heliospheric flux is from CMEs):

$$\frac{\pi(1AU)^2 (|B_{MAX}| - |B_{MIN}|) \cos(45^0)}{\overline{\varphi_0} (1 - \overline{D_0})} = \frac{1}{2} \left(\text{Coth} \left[\frac{\lambda}{2f_{MAX}} \right] - \text{Coth} \left[\frac{\lambda}{2f_{MIN}} \right] \right)$$

Estimate for disconnection rate

- Using:
 - $|B|_{MAX} = 9\text{nT}$ $|B|_{MIN} = 5\text{nT}$
 - $f_{MAX} = 3$ per day $f_{MIN} = 1/3$ per day
 - $D_0 = 0.4$ $\tau_0 = 10^{21}\text{Mx}$ (10^{20}Mx)
- $\tau = 0.7 \times 10^{-7} \text{ s}^{-1}$; i.e. $\tau \sim 12 \text{ days}$ (120 days)
 - If $V_{CME} \sim 400 \text{ km/s}$, this translates to $\sim 4 \text{ AU}$
- Too quick?

Lots of assumptions

- Heliospheric flux inferred from single point measurement
- All ICMEs are magnetic clouds
 - Selection effect: Fitted MCs contain more flux?
 - $\Phi_0 \sim 10^{20}$ Mx gives correct value for $\tau \sim 120$ days
- ICMEs can be represented by average values
- Observed ICME rates at min/max
- Observed rate of opening – limited dataset
 - Need more multi-point observations of ICMEs



Center for space physics
Boston University



Conclusions

- Flux contained in an MC can be significantly higher than force-free model estimates
 - However $\langle \rho_0 \rangle$ is probably over estimated
- Simple model of heliospheric flux, along with best parameter estimates, gives too fast a rate for ICME opening
 - Lots of room for maneuver
- Many assumptions and approximations made in making this estimate



Center for space physics
Boston University

