








Solar Origins of the October November 2003 Extreme Events



N. Gopalswamy

NASA/GSFC

*SHINE 2004 WG3 Thursday,
June 1 Big Sky, Montana*




In this Talk...

- Properties of the Solar Source
- Properties of the October – November CMEs in comparison with the general population of SOHO/LASCO CMEs
- 1-AU arrival times of CME-driven shocks
- Limits to CME speed and Maximum energy extractable from solar active regions

Extreme Solar Eruptions of October-November 2003


- Extreme flare size ($>X28$)
- Extreme CME speed (2700 km/s)
- Extreme concentration of energetic CMEs
- Extreme concentration of IP radio bursts
- Extreme AR area: 0.26% of solar surface
- Extreme SEP event: 33,000 pfu
- Extreme ICME magnetic field ~ 60 nT
- Extreme geoimpact: - 465 nT
- Extreme solar wind speed (~ 2000 km/s)

Eruption Centers




Recurrent Eruptions: Flares


- 143 flares from the three regions (X,M, & C only)
- Recurrence time of 3 h for dozens of flares.
- This means the regions released their free energy in fragments
- Average recurrence time is smallest for AR 484 (4.5 hours)
- Largest number of total, M and X flares from AR 486




At least
74 CMEs
From
3 ARs,
0484
0486
0488




2003/10/23 09:06




2003/10/23 20:30




2003/10/24 03:30




2003/10/24 05:54




2003/10/24 21:30




2003/10/25 05:30




2003/10/26 06:54




2003/10/28 11:30




2003/10/29 22:18



2003/11/01 23:30




2003/11/02 17:54





2003/11/04 20:06

Oct-Nov 2003 CMEs with respect to CMEs of cycle 23



Oct-Nov 03 Vs Cycle 23 CMEs



- Oct-Nov CMEs: faster & wider on the average



- The fastest CME (2657 km/s) occurred on Nov 04

- Large fraction of halo CMEs (13% Vs. 3%)

- Large fraction of ultrafast ($>2000 \text{ km/s}$) CMEs (11% Vs. 0.3%)

Extreme Concentration
Of energetic CMEs


Type II Radio Bursts: Shocks near the Sun





	UT	Metri	DH-	In
10/21	04:10	☐IV	km	situ
10/26	07:00	☐IV	☐	
10/26	17:45	☐IV	☐	☐
10/28	11:10	☐IV	☐☐	☐
10/29	21:15	☐IV	☐☐	☐
11/01	22:55	☐	☐	
11/02	09:23		☐☐	
11/02	17:30	☐IV	☐☐	☐
11/03	01:15	☐IV	☐	
11/03	10:00	☐IV	☐☐	
11/04	20:00	☐IV	☐☐	☐
11/05	01:40	x	☐	

12 DH type II bursts. 6 DH-km (~70 in cycle 23)
 11 WAVES IV bursts. (Only 12 other in cycle 23!)

Eight Shocks Reached Earth During the Oct-Nov Period




<http://umtof.umd.edu/pm/>




In situ Shocks & Associated CMEs

L1 Shock	CME (PA, speed, width)	Solar Region (Flare, AR, Loc)
10/24 14:47 UT	10/22 20:06 PA93 1085 km/s W=134	10/22 19:50 M9.9 AR0486 S22E90 ?
10/26 07:49 UT	10/23 08:54 PA088 1406 km/s W=115	10/23 08:19 X5.4 AR0486 S21E88
10/26 18:35 UT	10/23 20:06 PA101 1136 km/s W=089	10/23 19:50 X1.1 AR0486 S17E84
10/28 01:30 UT	10/26 17:54 PA 266 1537 km/s W=150	10/26 17:21 X1.2 AR 0484 S02W38
10/29 06:00 UT	10/28 11:30 PA092 2459 km/s W=360	10/28 09:51 X17.2 AR0486 S16E08
10/30 16:20 UT	10/29 20:54 PA092 2029 km/s W=360	10/29 20:37 X10.0 AR0486 S15W02
11/04 05:53 UT	11/02 17:30 PA262 2598 km/s W =360	11/02 17:03 X8.3 AR0486 S14W56
11/06 18:56 UT	11/04 19:54 PA257 2657 km/s W=360	11/04 19:29 X28 AR0486 S19W83


10/23 08:54




10/23 20:06




10/28 1054




10/29 20:37



11/02 17:30



11/04 19:54



October-November 2003: Solar Energetic Particle Events (> 1000 pfu)

$I_p = 10$ pfu (> 10 MeV) is considered hazardous.



Three $I_p > 1000$ pfu events occurred during Oct-Nov 03

(1 pfu = 1 proton/cm²/s/sr)

Only 33 such events since 1976

	Year	# of SEP Events
	1996	-
Each 1 denotes a >1000 pfu event	1997	-
	1998	1 1 Diff ARs
	1999	-
	2000	1 1 Diff ARs
	2001	1 1 1 1 1 Diff ARs
	2002	1 4/21
	2003	1 1 1 AR0486

All the three > 1000 pfu events (1) are from AR 486!




CMEs and SEPs

At least 5 large SEP Events

- Mostly from 486
- One from 0484
- 10/28 CME produced the largest > 10 MeV flux (33,600 pfu)


>10⁴ pfu Events: Cycle 23



10/28/03	33,600
11/04/01	31,700
07/14/00	24,000
11/22/01	18,900
11/08/00	14,800
09/24/01	12,900



Bigger (since 1976):

10/19/1989 – 40,000
03/21/1991 – 43,000



CME Speed vs 10 MeV Proton Intensity

- October Nov 2003 events extend the data points slightly to the right
- These plus the 1989 and 1991 events suggest an upper limit to the CME speed (~ 3000 km/s)




— Oct-Nov03
 — 89, 91

Diamond symbols
 Red: SEP events
 With pre-CMEs
 Blue: SEP events
 W/o pre-CMEs
 Green: Other
 Interactions
 (streamer interactio
 Below occulting disl

Shocks from 10/28-29 `03 CMEs Arrived at Earth in < 20 h

Flare onset to sudden commencement

Eruption Onset	TT (hrs)	Vcme (km/s)
04/15/38	21.2	2020
02/28/42	20.5	2080
02/07/46	17.8	2350
07/16/59	19.5	2170
08/04/72	14.6	2790
10/28/03	19.0	2459
10/29/03	19.7	2029
07/14/00	27.4	1670
10/26/03	31.5	1537




CME onset to shock arrival at L1

2790km/s

Fastest CME of Cycle 23


- 2003/11/04 CME: $V = 2700$ km/s
- Limb CME so speed is close to actual value
- Mass $\sim 2 \times 10^{16}$ g
- Kinetic Energy $\sim 7 \times 10^{32}$ erg
- The shock arrived after ~ 47 h because of grazing incidence.
- Similar to the inferred speed for the Aug 72 event



Active Region Area as a Fraction of the Solar Hemisphere

#	NOAA	AREA	DATE
01	0486	2610	2003/10/30
02	9393	2240	2001/03/29
03	9169	2140	2000/09/20
04	0069	1960	2002/08/19
05	0484	1750	2003/10/23
06	0488	1750	2003/10/31
07	0105	1520	2002/09/11
08	8323	1460	1998/09/04
09	9690	1420	2001/11/11
10	8651	1370	1999/08/03

About 0.6% of the disk was covered by ARs
 → Implications to irradiance



Total, Potential & Free Energies

- Potential $\frac{1}{8\pi} \int B_p^2 dV$ (from LOS B)
- Total $\frac{1}{8\pi} \int B_T^2 dV$ (from Vector B)
- Free Energy \sim Total – Potential
 \sim Potential Energy

See, however, Metcalf (2004), McTiernan (2004)


AR Potential Energy & CME Kinetic Energy

- AR 486: Volume $\sim 10^{31} \text{ cm}^3$
(300 x 300 x 300 arcsec³)
- Potential Energy $\sim 2 \times 10^{33} \text{ erg}$
- Total Magnetic Energy $\sim 2 \times \text{PE} \sim 4 \times 10^{33} \text{ erg}$
- Free energy $\sim 2 \times 10^{33} \text{ erg}$
- CME Kinetic Energy $\sim 7 \times 10^{32} \text{ erg}$
- $\sim 1/3$ of free energy goes into CME KE

For AR 7216 $U_t \sim 3.2 \times 10^{33} \text{ erg}$ (Metcalf et al. 1995)
All other forms of energy an order of magnitude lower
for large CMEs (Emslie et al. 2004 JGR)


Kinetic Energy of a 10^{17} g , 3000 km/s CME:
 $4.5 \times 10^{33} \text{ erg} \rightarrow$ limit of CME energy & Travel time

Potential Field Extrapolation From MDI Magnetogram




Courtesy: Yang Liu


CME Masses



CMEs Mass Distribution (SEP) Limb events (Up to 30 degrees off-limb)




$\langle M \rangle \sim 1.5 \times 10^{15}$ g for all CMEs
 (Vourlidas 2004)



Worst-Case Scenario

- There were only 25 of the 8000 CMEs had speed > 2000 km/s; only 4 with speed > 2500 km/s
- Inferred speeds of historical events is < 2800 km/s
- CMEs probably have a speed limit of ~ 3000 km/s
- This limit arises from the maximum energy extractable from an active region ($< 10^{34}$ erg)
- The Sun-Earth Travel time of shocks has a limit of \sim half a day



CMEs Don't Kill Dinosaurs


There is a limit to the maximum energy extractable from solar active regions

- Historical Shock Transit Times
- SOHO's Extensive Observations (M, V)
- Empirical Shock Arrival Model Curve
- Measured Size of Active Regions
- Measured Photospheric Magnetic Field (B)
- CME Kinetic Energy
- Estimated Magnetic Potential Energy


October-November 2003 Geomagnetic Storms

Storms of Cycle 23 with
Dst < -200 nT

- 00 2003/11/20 24:00 -
465
- 01 2003/10/30 23:00 -
401
- 02 2001/03/31 09:00 -
358
- 03 2000/07/16 01:00 -
301
- 04 2000/04/07 01:00 -
288
- 05 2001/11/06 06:00 -
277
- 06 2001/04/12 00:00 -
256
- 07 1999/10/22 07:00 -
237
- 08 2000/08/12 10:00 -
225






2003/10/28 11:30 2003/10/29 21:42 2003/11/02 17:54 2003/11/04 20:06



October-November 2003 Geomagnetic Storms: Return of the Dragon


Unusually Large Travel time for a fast CME. Why?

CME onset: 11/18 08:15
L1 shock: 11/20 07:30
Travel time ~ 47 h






Why did the 2003/11/18 CME-shock arrived so late?

V = 1660 km/s
T = 47 h





Extreme Deceleration of 11/18/03 CME


CORONAL HOLE DEFLECTS CME?



Extreme Impact on Geospace




Axial Fields of Cycle 23 MCs







Courtesy: N. Ness, R. Lepp

The Fastest CME of the cycle

Wind & SOHO





CME Interaction



Where is the Interaction Region?





- Nothing happening above the opposite limb
- The CME is at 18 Rs from the Sun just before interaction.
- The ambient density is expected to be 10^4 per cc.
- When the shock enters the dense structure, the upstream plasma frequency jumps \rightarrow frequency of emission jumps
- At the same time, the intensity is enhanced: Change in Alfvén speed
$$dV_a/V_a = dB/B - \frac{1}{2} dN/N$$
can strengthen the shock



Interaction detected by Wind, Ulysses & Cassini

Cassini at 8.7 AU:
Radiation has to travel
for additional 64 min

Wind: 21:10; Uly: 21:50; Cass: 22:18



Summary

- Record-setting speed (2657 km/s) on 2003/11/04 associated with the largest X-ray flare (X28); (previous: 2604 km/s on 5/12/2000)
- 2003 had 9 ultrafast (>2000 km/s) CMEs: 8 during Oct-Nov 2003 period. During cycle 23 (8 yrs) there were only 25 ultrafast CMEs. Only 4 CMEs had $V > 2500$ km/s
- Inferred speed of historical events is < 2800 km/s
- The Oct-Nov CMEs were faster and wider on the average \rightarrow SEP effective and Geoeffective
- 12 shocks near the Sun; 8 shocks detected *in situ*
- Both the Earth-directed CMEs produced intense geomagnetic storms (-363 & -401 nT). AR 501 (484) produced a CME on 11/18 that resulted in the largest storm of the cycle (-465 nT)
- About 1/3 of the magnetic free energy seems to go into the CME kinetic energy; Maximum free energy \sim a few 10^{33} erg
- There seems to be an upper limit to the CME speed ~ 3000 km/s, implying at least a half a day travel time for CME-driven shocks

Maximum Energy Extractable

- Free Energy = 5×10^{33} erg
- If all the energy goes into CMEs,


$$\frac{1}{2} MV^2 = 5 \times 10^{33} \text{ erg}$$

$$\Rightarrow V = 3160 \text{ km/s for } M = 10^{17} \text{ g}$$



Properties of Type IV Bursts

12 Events from 1997-2002

11 Events from 10/21-11/04 2003



1668 km/s



Lowest ending f
→ Tallest loops